



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

phyceae are the next younger group of algae, descended partly from Rhodophyceae and partly from flagellate-like organisms. (5) The Zygomycetes are derived from flagellated ancestors, the Peridinales being most nearly related to the modern flagellates. (6) The Chlorophyceae are the youngest of the algae, and have come partly from Rhodophyceae and partly from flagellated ancestors.—S. YAMANOCHI.

Sporangia and spores of *Aneimia*.—STEVENS²⁹ has investigated the development of the sporangia and spores in a species of *Aneimia*. He finds that the two tapetal layers break down at the mother cell stage, freeing the protoplasts and resulting in a tapetal plasmodium, as among the Ophioglossales. It was in connection with work on *Botrychium* (1906) that STEVENS proposed the excellent descriptive phrase "tapetal plasmodium." Perhaps it was a slip that he did not include this earlier paper in the "literature cited," or the still earlier paper of CARDIFF (1905). Upon the separation of the mother cells in *Aneimia* the plasmodium entirely surrounds each one. As each mother cell lies imbedded separately in the plasmodium, no wall is seen, and when the tetrad is formed the mother cell membrane persists about it. At the separation of the spores of a tetrad, the tapetal plasmodium flows between them. The author thinks that the thickness of the exine "is the work of the tapetal plasmodium." It is becoming more and more evident that in structure and behavior the Ophioglossales and Filicales belong together.—J. M. C.

Chromosomes in maize.—KUWADA³⁰ has studied the nuclear conditions in the pollen mother cells of nine different races of corn: red starch corn, yellow starch corn, amber rice popcorn, black starch corn, golden broach field corn, white flint corn, sugar corn, early light sugar corn, and red sugar corn. The number of gemini in these different races varies from 9 to 12, the sugar corns having generally a larger number than the starch corns. He thinks that the smaller number was reduced from 12, which is the original number for all the races of *Zea Mays*. The size and shape of the gemini in a figure differ, and there is present always a duplication of each of the gemini. In the equatorial plate of the homotypic division some pairs of chromosomes come in contact with each other. He suggests that the production of innumerable races of *Zea Mays* might have a certain relation to the duplication of chromosomes, resulting in the double number derived from the original form, which had probably 6 chromosomes as the reduced number.—S. YAMANOCHI.

***Botryopteris antiqua*.**—This interesting paleozoic fern, described by KIDSTON in 1908 from inadequate material, has been studied by Miss BENSON³¹

²⁹ STEVENS, WILLIAM CHASE, On the development of the sporangia and spores of *Aneimia phyllitidis*. Ann. Botany **25**:1059-1068. pls. 84, 85. 1911.

³⁰ KUWADA, Y., Maiosis in the pollen mother cells of *Zea Mays* L. Bot. Mag. Tokyo **25**:164-181. pl. 6. figs. 4. 1911.

³¹ BENSON, MARGARET, New observations on *Botryopteris antiqua* Kidston. Ann. Botany **25**:1045-1057. figs. 3. pls. 81-83. 1911.